



Effectiveness of a Physical Education Program on the Motor and Pre-literacy Skills of Preschoolers From the Training-To-Health Project: A Focus on Weight Status

Giuseppe Battaglia^{1,2}, Valerio Giustino^{1,2}, Garden Tabacchi^{1,2*}, Marianna Alesi¹, Claudia Galassi^{1,2}, Carmen Modica¹, Antonio Palma^{1,2} and Marianna Bellafiore^{1,2}

¹ Department of Psychological, Educational Science and Human Movement, University of Palermo, Palermo, Italy, ² Sport and Exercise Science Research Unit, University of Palermo, Palermo, Italy

OPEN ACCESS

Edited by:

Cristina Cortis,
University of Cassino, Italy

Reviewed by:

Stefano De Dominicis,
University of Copenhagen, Denmark
José A. Barela,
São Paulo State University, Campus
Rio Claro, Brazil

*Correspondence:

Garden Tabacchi
tabacchi.garden@libero.it;
garden.tabacchi@unipa.it

Specialty section:

This article was submitted to
Movement Science and Sport
Psychology,
a section of the journal
Frontiers in Sports and Active Living

Received: 02 July 2020

Accepted: 16 November 2020

Published: 16 December 2020

Citation:

Battaglia G, Giustino V, Tabacchi G,
Alesi M, Galassi C, Modica C,
Palma A and Bellafiore M (2020)
Effectiveness of a Physical Education
Program on the Motor and
Pre-literacy Skills of Preschoolers
From the Training-To-Health Project: A
Focus on Weight Status.
Front. Sports Act. Living 2:579421.
doi: 10.3389/fspor.2020.579421

Many studies reported a positive relationship between motor skills, cognitive functions, and school performance in school-age children; however, little is known in preschool children. The aim of the present study was to demonstrate the effectiveness of a physical education program (PEP) on locomotor, object control skills, and pre-literacy cognitive functions in a wide population of preschoolers and verify whether weight status could influence these abilities. In the context of the Training-to-Health Project, a sample of 1,029 preschoolers was recruited in kindergartens from the urban area of Palermo (Italy). Their gross motor and pre-literacy skills were tested before (PRE) and after (POST) following 16 weeks (2 h/week) of a PEP, which included ludic-motor activities aimed at developing body awareness and fundamental motor and perceptual-sensory skills. Analyses of variance (ANOVA) were performed to assess the skills before and after the intervention and to evaluate the effect of different categories of weight status on the examined variables. Regression analyses were conducted to confirm the hypothesized interrelationship between motor and pre-literacy skills in the considered sample. Both locomotor/object control and pre-literacy skills were significantly higher in children after the PEP ($p < 0.05$). We found 23% of overweight children and no significant difference between weight status classes in both PRE and POST PEP groups. In the POST group, higher locomotor and object control skills were mostly associated with better pre-literacy skills. This study shows that PEP was effective in improving both motor and pre-literacy skills in preschoolers independently from age and gender, while weight status did not affect these skills suggesting that this program can be administrated indifferently in children with different categories of weight status. Therefore, PEP could be a decisive education strategy to enhance motor and cognitive learning in preschool children and to achieve successful academic outcomes.

Keywords: physical education, motor skills, pre-literacy skills, preschooler, weight status

INTRODUCTION

Motor Skills

Fundamental movement skills (FMS), or simply motor skills, are defined as the basic level of movement and include locomotor skills (running, jumping, galloping, hopping, crawling) and object control (bouncing, catching, throwing, kicking) (Bellows et al., 2017; Han et al., 2018). The concept of FMS is strictly related to the quality of motor skill competence (MC), which is the level ability of motor coordination, and it is measured through several test batteries (D'Hondt et al., 2009; Lopes et al., 2012; Figueroa and An, 2017; Maeng et al., 2017; Loras, 2020; Tsuda et al., 2020).

The appropriate development of FMS, which occurs during the preschool period (children aged 3–5 years), allows to build the basis for performing complex movements and task-specific abilities (Figueroa and An, 2017; Aivazidis et al., 2019; Alves and Alves, 2019). Moreover, several studies reported that the practice of physical activity (PA) improves FMS level by inducing a higher MC in childhood (Fisher et al., 2005; Hardy et al., 2010; Battaglia et al., 2018; Aivazidis et al., 2019; Nilsen et al., 2020a). For these reasons, the practice of structured PA in kindergarten during this period results to be crucial (Deli et al., 2006; Barbosa et al., 2016; Figueroa and An, 2017; Kippe and Lagestad, 2018; Alves and Alves, 2019; Jarraya et al., 2019). However, PA in preschool age is limited and, furthermore, the level of PA among children living in Europe varied considerably among countries (Konstabel et al., 2014). The latter topic is also confirmed by the fact that over the years several research groups have focused on the importance of PA in children and adolescents, and the few articles present in the literature about this issue in preschooler demonstrate the recent interest among scientists (Van Capelle et al., 2017; Engel et al., 2018).

It is well-known that PA plays a key role in promoting health and preventing diseases at every stage of life, including children and adolescents (Nielsen et al., 2016; Voss and Harris, 2017; Czenczek-Lewandowska et al., 2018). The lack/low level of PA practice and sedentary behaviors negatively affect body weight in preschooler children (Han et al., 2010; Osinski and Kantanista, 2017; Rodriguez-Ayllon et al., 2019). Furthermore, the assumption of harmful behaviors (e.g., eating unhealthy foods) during the time spent in a sedentary lifestyle has been shown to exacerbate the capacity to maintain an adequate body weight in children by creating a vicious cycle (Pagani et al., 2010; Eichinger et al., 2017). As reported by several researches, both physical inactivity and sedentary/unhealthy behaviors increase the risk of being overweight and obesity in childhood (Tucker, 2008; Han et al., 2010; Hills et al., 2011; Reilly et al., 2019). For this reason, the role of diet and physical activity in children has been investigated in previous studies (Obradovic Salcin et al., 2019; Tabacchi et al., 2020a). In support of the above, Tabacchi et al. (2020a) suggest that raising children in an environment where both motor and cognitive skills are encouraged can improve food literacy and, furthermore, enhance school achievement.

The prevalence rate of overweight and obesity among European children, with different rates between countries, is about 20% and represents a public health problem due to the

related risk factors (Ahrens et al., 2014; Eichinger et al., 2017). In order to contrast this phenomenon, a series of political and social initiatives and experimental approaches have been promoted by encouraging the practice of PA among preschoolers (Palmer et al., 2019; Hoza et al., 2020; Popovic et al., 2020).

The beneficial effects of PA for preventing overweight and obesity are well-documented. Furthermore, as reported in the review by Bingham et al. (2016), the interrelationship between PA and several socio-ecological domains, such as anthropometric/demographic features (e.g., age, gender, weight status), and family community level/parental behaviors (e.g., family socioeconomic status/parents' PA level) is widely recognized in early children (Kimbro et al., 2011; Eichinger et al., 2017). In a similar way, FMS have reported to positively affect psychosocial aspects and cognitive functions in preschool-aged children (Rhemtulla and Tucker-Drob, 2011; Oberer et al., 2017). However, only a few studies have investigated the influence of motor skills on PA levels and preschoolers' body weight status, reporting conflicting outcomes (D'Hondt et al., 2009; Logan et al., 2011; Morano et al., 2011; Lopes et al., 2012; Roscoe et al., 2019).

Pre-literacy Skills

Positive influence of PA programs on pre-literacy skills has been found in the kindergarten age (Barnett et al., 2008; Callcott et al., 2015; Mavilidi et al., 2015, 2017).

Pre-literacy refers to kindergarten skills that are predictors of later literacy achievement. These skills include a set of competences, such as (1) knowledge and understanding about printed materials; (2) oral language abilities, e.g., vocabulary, comprehension, and listening; and (3) alphabetic code awareness, e.g., phonological/phonemic abilities to detect and manipulate syllables, phonemes, and word parts (Lonigan and Shanahan, 2010; Puranik and Lonigan, 2011; Pinto et al., 2016).

Given these positive influences of PA on pre-reading/writing and pre-math, researchers and practitioners have developed a variety of preschool-based programs suitable to enhance pre-literacy skills through PA activities. The underlying conceptualization is that active play characterizing preschool-age PA would be a natural and enjoyable context to encourage linguistic development by creating more frequent linguistic and social opportunities, to understand and explain instructions or play rules, to elaborate stimuli from many sources, to experiment direct or indirect performances, and to try new action-based representations of tasks (Boncoddio et al., 2010; Carson et al., 2015).

For example, the Movement with Literacy (MowL) was found to improve phonological awareness, spelling, and motor coordination. It was a program based on a 15-min daily preschool curriculum composed of literacy (Let's Decode) and movement (Moving on with Literacy) curricula (Callcott et al., 2015) and counted in 30 action songs with motor tasks to train fine and gross motor, eye-tracking, balance, rhythm, core strength, and aerobic skills. Another program delivered by classroom teachers over 8 months was made up of 60-min moderate PA units (2 times per day) associating motor (jumping, running,

moving on lines, marching) and early literacy tasks to train oral language, vocabulary, and phonological awareness (Kirk et al., 2014; Kirk and Kirk, 2016). Programs implemented by Mavilidi et al. (2015, 2017) improved phonological and science competences in preschool age. Rhyming, alliteration, and picture naming skills were enhanced through PA activities that associated full-body movements, e.g., physical exercises, or part-body movements, e.g., gestures, and correspondent foreign words. Science competences were improved by an integrated method that trained children to learn planets' names and their correct position from the sun by performing movements from the sun to the closest planet and so on.

Moreover, a Parent-oriented Movement and Pre-literacy Program required 60 min a week over 10 weeks of activities and involved both preschool age children and their parents. It consisted of Fundamental Movement tasks; free-play activities with balls, steps, bricks, or puzzles; and a storybook reading activity shared among children and their parents to enhance motor and literacy skills as print-concept and alphabet knowledge (Bedard et al., 2017).

Purpose

In our previous work we investigated the effect of a 16-weeks physical education program (PEP) in motor and cognitive preschool children's status, finding an improvement on both domains (i.e., motor skills and cognitive functions) (Battaglia et al., 2018). However, as stated in that paper, in the present study we applied the further step for the validation of the study, i.e., to extend the investigation to a larger sample size in order to illustrate the effects of the developed PEP on locomotor, object control skills, and pre-literacy cognitive functions in a larger scale (Battaglia et al., 2018). Moreover, based on this large preschooler sample, the further purpose of the study was to investigate whether weight status could influence these skills.

MATERIALS AND METHODS

Study Design and Participants

The present study is a non-randomized trial carried out within the Training-to-Health Project, financed by the Municipality of Palermo in 2016 with the general aim of enhancing motor and cognitive skills in preschoolers, beyond other scopes, such as monitoring aspects related to these abilities (Tabacchi et al., 2020b). A team of experts from the University conducted the study and recruited physical education specialists (PESs) with previous experience in the field of motor and cognitive science in children, in order to carry out activities within the selected kindergartens. Teachers from the classes were directly involved as support in the program activities. All the personnel taking part in the program was properly trained, and methodologies were standardized to allow the collection of accurate and reliable data.

The study was approved by the Ethical Board of the University of Palermo (N. 2/2018) and followed the criteria for the use of persons in research as defined in the Declaration of Helsinki.

The preschoolers' sampling envisaged a mixed multistage procedure. In the first stage, all the kindergartens of Palermo city boundaries (Municipality of Palermo) ($n = 389$) were identified;

in the second stage, the sample was stratified into public ($n = 193$) and private kindergartens ($n = 196$) and the cluster of public kindergartens was selected; in the third stage, $n = 21$ (11%) public kindergartens were randomly chosen in geographical areas stratified by different socioeconomic environments (SEE), every class within the chosen cluster was chosen, and every student within the class was sampled ($n = 1,054$). The SEE was denoted according to the "index of socio-economic disadvantage," measured on the basis of four indicators of deprivation in the 55 different city districts (http://cqd.comune.palermo.it/CQDSupera/docs/CQD4982_1221223025117.pdf).

Parents were asked to sign an informed consent to let their children participate the study. Among them, a total of 25 refused to sign (2.4%), and a final sample of 1,029 children was obtained.

Since collecting measures for all the individuals was not possible, as some of them were not present the day of the assessment of a particular item, some results are referred to a slightly lower sample, as indicated in the result tables.

Study Procedure

Participants' weight and height were initially measured, and they were assessed for basic motor skills and pre-literacy abilities by the PESs (we will further indicate this group as the "PRE group").

The motor skill items were measured by the Italian version of the gross motor development test (Ulrich, 2003), which consists in two different aspects of gross motor development, i.e., locomotion (requiring subjects to run as fast as possible for 15 m, jump forward, gallop for 10 m, hop on one leg for 5 m, do a long jump, and take little jumps forward and laterally), and object control (bounce the ball, catch the ball, catch a ball with a tennis racket, and running while kicking a ball and throwing a ball). The combination of these two subtests provides the quotient of gross motor development (QGMD), useful for assessing the overall gross motor skills of children.

Pre-literacy abilities were assessed by choosing four tasks related to visual analysis and spatial orientation abilities, derived from the Italian battery PRCR-2/2009 (Cornoldi et al., 2009), measuring general and specific prerequisites to reading and writing abilities in preschoolers.

Details of these assessment tools are described in our previous paper (Battaglia et al., 2018).

After the initial assessment, preschoolers were subjected to a Physical Education Program (PEP) for 16 weeks, applied with a frequency of twice a week, by the PESs. In agreement with Battaglia's PEP (Battaglia et al., 2018), each lesson (~60 min) included a warm-up and social interaction phase (~5 min) which aimed to enhance the fitness level of children and their motivation to participate in activities (e.g., circle time in which children, sitting on the floor, greeted everyone and took turns performing a movement requested by the physical education specialist; activities of running, jumping, catching/throwing/kicking a ball, etc.); a central phase (~50 min) including specific activities in order to develop perceptual-sensory and fundamental motor and skills in preschool children; and a cool-down phase (~5 min) in order to relax children and explore their level of satisfaction (e.g., circle time in which children, lying on the ground, performed calming breathing

activities). At the end of the PEP, preschoolers were assessed again for the gross motor and pre-literacy skills, and also anthropometric measures were collected again. This group following the PEP represents the "POST group." Children of the PRE group were included in classroom activities for an equal time as the POST group with teachers. Both groups completed the activities during the school period in a multi-activity area.

BMI and Motor Skill Measures

Age of the preschoolers was retrieved from the birth dates reported in the kindergartens' lists.

Weight and height were measured through Seca electronic scales (maximum weight recordable, 300 kg; resolution, 100 g; Seca Deutschland, Hamburg, Germany) and stadiometers (maximum height recordable, 220 cm; resolution, 1 mm) which were used according to standard procedures (Lohman et al., 1988). Body mass index (BMI) was calculated, and four classes of weight status were obtained according to the BMI percentiles for males and females aged 0–18 years from Cole et al. (2000): underweight below the 5th percentile, normal weight between the 5th and 85th percentiles, overweight between the 85th and 95th percentiles, and obese over the 95th percentile.

The gross motor skills were measured as scores derived from the marks attributed to the single performance. All participants completed three trials of each gross-motor skill and acquired a "1" mark, when a criterion was used incorrectly two out of three times or was not observed, or a "0" grade, when a criterion performance was executed two out of three times. According to the age level of the participant, the sum of the scores detected for each item (maximum total score 48) was converted into standard scores. The QMGD was obtained by summing up the scores of the two subtests for assessing locomotor and object control skills, and it could score from 46 to 154.

The categorization of QGMD in classes of motor abilities was performed according to the manual (Ulrich, 2003): 35–69 (very low motor ability, VL-MA), 70–79 (low motor ability, L-MA), 80–89 (below average motor ability, UA-MA), 90–110 (average motor ability, A-MA), 111–120 (over average motor ability, OA-MA), 121–130 (high motor ability, H-MA), and 131–165 (very high motor ability, VH-MA).

Pre-literacy Skill Measures

Pre-literacy skills were measured through four tasks derived by PRCR-2/2009 (Cornoldi et al., 2009) which is an Italian battery of standardized tasks to test kindergarten children's prerequisites to later reading and writing abilities. The tasks were (1) printed-letter identification aimed at measuring visual analysis and spatial orientation abilities; (2) object naming aimed at measuring linguistic proficiency, the visual attention, and the sequentiality of eye movements; (3) partially hidden-object naming aimed at measuring linguistic proficiency, the visual attention and discrimination, and the sequentiality of eye movements; and (4) pointed-object naming aimed at measuring the visuo-perceptual ability to identify a figure from the background, the linguistic proficiency, the visual attention and discrimination, and the sequentiality of eye movements. More in detail, the printed-letter identification task consisted of a white sheet with 12 target letters

printed on the left and four letters for each target (the target and three distracting letters) printed on the right. Each child was asked to identify and cross the letter corresponding to the right target. The object-naming task was composed of 30 objects in five sequences of six objects for each (e.g., animals, flowers, ice cream, sun, star), and each child was asked to name each object. The partially-hidden-object naming task was composed of three sequences of objects, already shown in the object-naming task, but the objects were overlapping and smaller. Each child was asked to recognize and name each object. The pointed-object-naming task was composed of two sequences of overlapping objects that appeared in the partially hidden-object-naming task with four objects for each sequence marked by a dot at 15 mm. Each child was asked to rapidly name the marked objects from left to right and from top to bottom.

The criteria of evaluation were the sum of errors and the time of performance.

Statistical Analysis

Sample descriptive data are shown in number and percentages for categorical outcomes and mean and Standard Deviation (SD) for normally distributed continuous data. Normality of data was evaluated through the skewness/kurtosis test for normality.

The difference between PRE and POST groups skills was calculated through analysis of variance (ANOVA) adjusted for age and gender. This analysis was also used to assess differences in gross motor skills and pre-literacy abilities by classes of weight status.

Linear regression analyses were conducted to assess the degree of correlation between gross motor and pre-literacy abilities. Significance was set at $p < 0.05$. The software STATA.12 was used to perform the statistical analyses.

RESULTS

Sample Characteristics

The characteristics of the sample are shown in **Table 1**.

Differences Between PRE and POST PEP

The analysis of variance showed that the QGMD was significantly higher in the POST group compared to the PRE group (mean 108.8 vs. 123.9, $p < 0.001$); according to the QGMD classes, it passes from "average motor ability" in the PRE group to "high motor ability" in the POST group. This high difference was present also for the components of the QGMD, i.e., locomotor (mean 10.7 vs. 14.0) and object control skills (mean 12.3 vs. 14.1) (**Table 2**). Moreover, all the evaluated locomotor and object control subitems were significantly higher in the test group (**Table 2**), with an average score increase of 0.63 units (range 0.5–0.8) and 0.55 units (range 0.47–0.7), respectively.

With regard to the pre-literacy skills, all of them significantly increased in the POST PEP group, meaning that both errors and time needed to perform the test were significantly lower in this group compared to the PRE group (**Table 3**). For example, the errors in the identification of printed letters were on average 3.4 in the PRE group and they decreased to 2.7 in the POST group; the time needed to naming objects

TABLE 1 | Characteristics of the sample before (PRE) and after (POST) the physical education program.

	No.	%		
Gender				
Males	557	54.1		
Females	472	45.9		
Tot	1,029	100		
Age				
	Pre		Post	
	No.	%	No.	%
3 years (≤ 47 months)	200	19.4	142	13.8
4 years (48–59 months)	393	38.2	380	36.9
5 years (≥ 60 months)	436	42.4	507	49.3
Tot	1,029	100	1,029	100
Weight status				
Normal	587	69.5	555	66.5
Under	90	10.7	87	10.4
Over	127	15.0	130	15.6
Obese	41	4.9	62	7.4
Tot	845	100	834	100
Mean and SD				
	Pre		Post	
	Mean	SD	Mean	SD
Age (months)	56.11	9.85	58.98	10.03
Height (m)	1.08	0.07	1.09	0.08
Weight (kg)	19.05	3.93	19.36	4.13
BMI (kg/m^2)	15.94	2.02	16.07	2.31

decreased from 72.8 to 62.2 s. On average, the number of errors decreased by 1.05 units (range from 0.7 to 2 units), while the time needed in naming objects decreased of 14.5 s (10.6 and 18.4 s in naming objects and naming partially hidden objects, respectively).

Differences by Weight Classes

When analysis of variance was conducted by categories of weight status, the QMGD, locomotor, and object control skills were not significantly different throughout the classes of underweight, normal overweight, and obesity (Table 4). Actually, observing all the skill values in obese preschoolers, reduced performances can be evidenced, even though they were not statistically significant. The same observation can be done for the pre-literacy skills, with a trend of decreasing performances from normal weight to obese individuals (Table 5) but no statistical significance in those differences. Similar results were found out also for all the locomotor subitems, with the exception of leaping ($p < 0.05$), and for all the object control subitems (Supplementary Material 1).

These results were found out both in POST and PRE PEP groups (PRE group results are not shown in the present paper).

Regressions of Motor and Pre-literacy Skills (POST)—Adjusted for Age and Gender

The regression analysis showed that children with high QGMD make less errors in naming objects and partially hidden objects (coefficients -1.22 and -0.49) and take less time in naming pointed objects (coefficients -0.55) (Table 6). The component of object control skills is significantly correlated with all the pre-literacy skills, with higher object control ability associated with better pre-literacy ability (coefficients ranging from -0.04 to -1.13); for the locomotor skills, the time needed in performances was not correlated, but all the other items (errors) were with coefficients ranging from -0.16 to -0.86 (Table 6).

These results are confirmed when analyzing the locomotor and object control subitems (Supplementary Material 2).

DISCUSSION

The aim of the present study was to investigate the effect of a physical education program (PEP) on motor and cognitive skills in a large preschooler sample focusing on the influence of children's weight status.

As we hypothesized, our findings showed an improvement of the QGMD level and in both of its components, i.e., locomotor and object control skills, after the PEP. These results are in agreement with the outcomes we found in our previous pilot trial in which we examined the effectiveness of this PEP on the aforementioned skills in a sample of 119 preschool children (Battaglia et al., 2018). Since, as we reported in that work, the main issue was to extend the PEP on a larger scale in order to validate its effectiveness, the present study allowed us to confirm previous results.

Our findings are consistent with scientific evidence from the literature demonstrating the positive effect of PA on improving motor and pre-literacy skills in preschool age (Orton et al., 2009; Logan et al., 2012; Zeng et al., 2017; Aivazidis et al., 2019; Popovic et al., 2020). Indeed, although the literature regarding the practice of PA in kindergarten is limited, recent researches have investigated the effects of different PA interventions on several skills and health aspects in preschoolers (Popovic et al., 2020; Toussaint et al., 2020). In 2008, Stodden et al. developed a conceptual model hypothesizing a primary connection between the level of PA and the quality of MC (Stodden et al., 2008). The findings of this seminal work have been confirmed by several subsequent research groups (Lai et al., 2014; Engel et al., 2018; Nilsen et al., 2020b; Xin et al., 2020). Goodway et al. (2003) showed the positive effects of a 9-weeks integrative PA program (18 lessons, 35/min each) on the Test of Gross Motor Development (TGMD) scores in preschoolers compared to children who only carried out the regular kindergarten PA. Based on these results, the authors supported the idea of implementing PA in the kindergarten education program (Goodway et al., 2003). Likewise, DuBose et al. (2018) found a positive relationship between the level of PA in children, measured through an accelerometer, and motor skills. In the same way, pre-literacy skills in preschoolers are positively related to PA

TABLE 2 | Evaluation of the locomotor and object control skills before and after the physical education program.

	PRE		POST		N	p-Value*
	Mean	SE	Mean	SE		
QGM	108.8	0.78	123.9	0.62	805	0.0000
Locomotor skills	10.7	0.17	14.0	0.15	807	0.0000
Locomotor subitems						0.0000
Running	2.6	0.04	3.2	0.04	966	0.0000
Galloping	2.0	0.05	2.7	0.05	967	0.0000
Hopping	1.7	0.05	2.5	0.05	967	0.0000
Leaping	1.2	0.04	1.7	0.04	967	0.0000
Horizontal jumping	2.2	0.04	2.9	0.05	967	0.0000
Skipping	1.3	0.04	1.8	0.04	967	0.0000
Sliding	2.3	0.05	2.9	0.05	967	0.0000
Object control skills	12.3	0.14	14.1	0.15	966	0.0000
Object control subitems						
Two-hand striking	1.4	0.04	2.1	0.05	967	0.0000
Stationary bouncing	1.03	0.03	1.5	0.04	967	0.0000
Catching	2.2	0.04	2.8	0.05	967	0.0000
Kicking	2.0	0.04	2.5	0.05	967	0.0000
Overhand throwing	1.9	0.05	2.4	0.05	966	0.0000

QGM, Quotient of Gross Motor Development.

*The difference between pre and post was calculated through ANOVA adjusted for age and gender. Significance set at $p < 0.05$.

TABLE 3 | Evaluation of the pre-literacy skills before and after the physical education program.

	PRE		POST		N	p-Value*
	Mean	SE	Mean	SE		
Printed-letter identification (error n.)	3.4	0.10	2.7	0.10	751	0.0000
Object naming (s)	72.8	1.09	62.2	0.89	713	0.0000
Object naming (error n.)	2.0	0.09	1.2	0.07	709	0.0000
Partially-hidden-object naming (s)	116.6	1.85	98.2	1.37	688	0.0000
Partially-hidden-object naming (error n.)	5.6	0.20	3.6	0.16	730	0.0000
Pointed-object naming (error n.)	2.3	0.06	1.6	0.06	698	0.0000

*The difference between pre and post was calculated through ANOVA adjusted for age, gender. Significance set at $p < 0.05$.

TABLE 4 | Differences in QGM, locomotor, and object control skills by categories of weight status.

	QGM			Locomotor skills			Object control skills		
	Mean	SD	p-Value	Mean	SD	p-Value	Mean	SD	p-Value
Weight status			0.0843			0.3247			0.0578
Normal	123.0	17.22		13.5	3.73		14.3	2.86	
Under	125.6	16.92		13.8	3.43		14.7	2.71	
Over	124.3	18.46		13.8	3.79		14.4	3.20	
Obese	117.2	19.93		12.6	3.91		13.2	2.41	

programs (Carson et al., 2016; Donnelly et al., 2016; Zeng et al., 2017).

As abovementioned, the results we found regarding the effect of PA on motor and pre-literacy skills are in line with several authors (Callcott et al., 2015; Mavilidi et al., 2015, 2017; Kirk

and Kirk, 2016; Bedard et al., 2017). Children with high QGM performed better on PRCR-2 tasks. These results revealed in these children a higher level on the abilities of visual analysis, visual attention, visual discrimination, spatial orientation, and linguistic proficiency. This could be due to PEP that should

TABLE 5 | Differences in pre-literacy skills by categories of weight status.

	Printed-letter identification (error n.)			Object naming (s)			Object naming (error n.)		
	Mean	SD	<i>p</i> -Value	Mean	SD	<i>p</i> -Value	Mean	SD	<i>p</i> -Value
(A) Printed letters identification and objects naming									
Weight status			0.1593			0.1078			0.6410
Normal	3.0	2.98		63.0	23.89		1.3	1.06	
Under	2.36	2.32		68.4	28.23		1.1	0.63	
Over	2.55	1.68		62.9	27.73		1.1	0.63	
Obese	1.75	1.58		55.0	25.35		0.9	0.68	
	Partially-hidden-object naming (s)			Partially-hidden-object naming (error n.)			Pointed-object naming (error n.)		
	Mean	SD	<i>p</i> -Value	Mean	SD	<i>p</i> -Value	Mean	SD	<i>p</i> -Value
(B) Partially-hidden-object naming and pointed-object naming									
Weight status			0.3365			0.7230			0.9670
Normal	101.4	36.53		3.8	4.67		1.61	1.45	
Under	107.6	45.72		3.4	3.28		1.6	1.47	
Over	100.6	32.93		3.5	4.85		1.63	1.56	
Obese	90.2	39.79		2.9	3.78		1.56	1.55	

TABLE 6 | Results of the regression analysis between motor and pre-literacy skills.

	QMGD				Locomotor skills				Object-control skills			
	Coef	SE	<i>p</i>	<i>N</i>	Coef	SE	<i>p</i>	<i>N</i>	Coef	SE	<i>p</i>	<i>N</i>
Printed letters	-0.25	0.248	0.324	707	-0.28	0.087	<i>0.001</i>	743	-0.18	0.064	<i>0.005</i>	744
Object naming time (s)	-0.03	0.031	0.355	673	-0.04	0.011	<i>0.000</i>	708	-0.004	0.008	<i>0.576</i>	709
Object-naming errors (n.)	-1.22	0.377	<i>0.001</i>	666	-0.59	0.129	<i>0.000</i>	702	-0.32	0.093	<i>0.001</i>	702
Partially-hidden-object-naming time (s)	-0.03	0.019	0.136	666	-0.02	0.007	<i>0.008</i>	701	-0.001	0.005	0.818	701
Partially-hidden-object-naming errors (n.)	-0.49	0.138	<i>0.000</i>	698	-0.25	0.048	<i>0.000</i>	734	-0.16	0.035	<i>0.000</i>	734
Pointed-object naming	-0.55	0.453	<i>0.000</i>	679	-1.13	0.157	<i>0.000</i>	715	-0.86	0.114	<i>0.000</i>	715

The italic was used to underline the significant values ($p < 0.05$).

enable kindergarten children to train visuospatial abilities, which are necessary to master print directionality, to align or put in a column the numbers with consequent accuracy on computation and numerical representations. Visual-motor integration abilities are essential to organizing and spacing letters, words, and numbers on a page. Moreover, motor tasks as jumping, running, moving on lines, and marching revealed to be suitable to train the ability to recognize and produce alliteration and rhyming (Mavilidi et al., 2015, 2017; Kirk and Kirk, 2016). In the context of interaction with peers and movement exercises based on rhythm, fine and gross motor practice, eye-tracking, and balance, children have more opportunities to develop their linguistic and metalinguistic skills, such as alphabet knowledge, letter-sound knowledge, detection and manipulation of phonemes or syllables, and the knowledge to encode and decode words.

While when the effects of PA were analyzed by weight status categories, no significant differences were found on the QMGD and locomotor and object control skills between underweight, normal overweight, and obesity, in both PRE and POST PEP groups. Although few studies have investigated

the relationship between motor skills and weight status/BMI domain in preschool age, the findings are conflicting (Lopes et al., 2012; Kim and Lee, 2016; Augustijn et al., 2018; Roscoe et al., 2019). For instance, D'Hondt et al. (2009) showed significantly higher levels of balance, object control, and manual dexterity in children with normal weight and overweight than obese peers. Similarly, Logan et al. (2011) reported significant differences on MC among children ages 4–6 years with a different BMI status indicating lower scores of motor skill proficiency in overweight and obese children compared to normal-weight peers. In contrast, the results reported by a large body of increasing research showed no correlation between FMS and weight status/BMI levels in preschoolers, as we found (Saraiva et al., 2013; Kim and Lee, 2016; Mülazimoglu Balli, 2016).

The key explanation could lie in the fact that cortical areas reach different levels of maturity of each other during the development process, and, moreover, the relationship between executive functioning and weight status in children has only been discovered in some cerebral areas (Liang et al., 2014). Among the

latter, the literature suggests that at an early age the prefrontal cortex results are less developed and this physiological condition does not allow to identify any differences in motor and pre-literacy skills among preschoolers belonging to distinct weight classes (Liang et al., 2014).

Although we found no difference between weight status groups, our results showed lower scores in leaping in children classified as overweight/obese. We suppose that weight status adversely affects this skill performance. Our hypothesis is sustained by several researches in which it has been demonstrated that tasks requiring propulsion are strictly dependent on body mass, reporting an inverse relationship between the amount of body mass and the lower-limb strength scores, which is an association between motor skills and overweight/obese status only for some locomotor skills (Siahkoughian et al., 2011; Castetbon and Andreyeva, 2012).

Therefore, our outcomes seem to suggest that this PEP can be administered in preschoolers with different weight status categories likewise, in order to improve motor and cognitive skills and to provide favorable bases for enabling positive academic achievement.

In the conceptual model of Stodden et al. (2008), an association between higher level of PA and greater scores in both locomotor and object control skills has been suggested. Moreover, the authors support the notion that preschool children with poor level of FMS could be more physically inactive in the future, leading, among other things, to an increased risk of being overweight/obese (Stodden et al., 2008).

As preschoolers currently perform lower than recommended PA levels by increasing the associated health risk and compromising skill development and later school achievement, we suggest that, based on our results, children would spend more time in structured PA in kindergartens (Gu, 2016; Kippe and Lagestad, 2018; Kobel et al., 2019). Therefore, we suggest that this PEP could be adopted as an educational strategy to improve motor and pre-literacy skills in preschool children, known to be a peculiar growth period for the development of these characteristics. Furthermore, we suggest that the activities of this PEP can also be proposed by teachers online, a modality that could be useful in critical and peculiar periods in which the level of physical activity could decrease as during the current COVID-19 pandemic (An, 2020; Giustino et al., 2020). To sum up, the implementation of enjoyable methods of teaching to

enhance children's school readiness through motor activities is an important matter to be investigated by educational scientists.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Ethical Board of the Azienda Ospedaliera Universitaria Policlinico Paolo Giaccone Palermo (Palermo 1, N. 02/2018). Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

AUTHOR CONTRIBUTIONS

MB conceptualized the paper. CG and CM collected the data. GT carried out the formal curation and analysis of data and drafted the results. GB, VG, and GT drafted original draft. MB, MA, and GB reviewed and edited the manuscript. AP supervised the final manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This research was funded by the Municipality of Palermo (grant number PJ_CT_D12_EDUSAN_0001).

ACKNOWLEDGMENTS

The authors wish to thank city councilor Barbara Evola, municipal manager Maria Anna Fiasconaro, schoolteachers, and physical education specialists for their kind collaboration in this study.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fspor.2020.579421/full#supplementary-material>

REFERENCES

- Ahrens, W., Pigeot, I., Pohlmann, H., De Henauw, S., Lissner, L., Molnar, D., et al. (2014). Prevalence of overweight and obesity in European children below the age of 10. *Int. J. Obes.* 38, 99–107. doi: 10.1038/ijo.2014.140
- Aivazidis, D., Venetsanou, F., Aggeloussis, N., Gourgoulis, V., and Kambas, A. (2019). Enhancing motor competence and physical activity in Kindergarten. *J. Phys. Act. Health* 16, 184–190. doi: 10.1123/jpah.2018-0260
- Alves, J. G. B., and Alves, G. V. (2019). Effects of physical activity on children's growth. *J. Pediatr.* 95, 72–78. doi: 10.1016/j.jpmed.2018.11.003
- An, R. (2020). Projecting the impact of the coronavirus disease-2019 pandemic on childhood obesity in the United States: a microsimulation model. *J. Sport Health Sci.* 9, 302–312. doi: 10.1016/j.jshs.2020.05.006
- Augustijn, M., D'Hondt, E., Van Acker, L., De Guchteneere, A., Lenoir, M., Caeyenberghs, K., et al. (2018). Role of motor competence and executive functioning in weight loss: a study in children with obesity. *J. Dev. Behav. Pediatr.* 39, 642–651. doi: 10.1097/DBP.0000000000000589
- Barbosa, S. C., Coledam, D. H., Stabelini Neto, A., Elias, R. G., and Oliveira, A. R. (2016). School environment, sedentary behavior and physical activity in preschool children. *Rev. Paul. Pediatr.* 34, 301–308. doi: 10.1016/j.rpped.2016.01.001
- Barnett, L. M., Van Beurden, E., Morgan, P. J., Brooks, L. O., and Beard, J. R. (2008). Does childhood motor skill proficiency predict adolescent fitness? *Med. Sci. Sports Exerc.* 40, 2137–2144. doi: 10.1249/MSS.0b013e31818160d3
- Battaglia, G., Alesi, M., Tabacchi, G., Palma, A., and Bellafiore, M. (2018). The development of motor and pre-literacy skills by a physical education program

- in preschool children: a non-randomized pilot trial. *Front. Psychol.* 9:2694. doi: 10.3389/fpsyg.2018.02694
- Bedard, C., Bremer, E., Campbell, W., and Cairney, J. (2017). Evaluation of a direct-instruction intervention to improve movement and preliteracy skills among young children: a within-subject repeated-measures design. *Front. Pediatr.* 5:298. doi: 10.3389/fped.2017.00298
- Bellows, L. L., Davies, P. L., Courtney, J. B., Gavin, W. J., Johnson, S. L., and Boles, R. E. (2017). Motor skill development in low-income, at-risk preschoolers: a community-based longitudinal intervention study. *J. Sci. Med. Sport* 20, 997–1002. doi: 10.1016/j.jsams.2017.04.003
- Bingham, D. D., Costa, S., Hinkley, T., Shire, K. A., Clemes, S. A., and Barber, S. E. (2016). Physical activity during the early years: a systematic review of correlates and determinants. *Am. J. Prev. Med.* 51, 384–402. doi: 10.1016/j.amepre.2016.04.022
- Boncoddo, R., Dixon, J. A., and Kelley, E. (2010). The emergence of a novel representation from action: evidence from preschoolers. *Dev. Sci.* 13, 370–377. doi: 10.1111/j.1467-7687.2009.00905.x
- Callcott, D., Hammond, L., and Hill, S. (2015). The synergistic effect of teaching a combined explicit movement and phonological awareness program to preschool aged students. *Early Childhood Educ. J.* 43, 201–211. doi: 10.1007/s10643-014-0652-7
- Carson, V., Hunter, S., Kuzik, N., Wiebe, S. A., Spence, J. C., Friedman, A., et al. (2016). Systematic review of physical activity and cognitive development in early childhood. *J. Sci. Med. Sport* 19, 573–578. doi: 10.1016/j.jsams.2015.07.011
- Carson, V., Kuzik, N., Hunter, S., Wiebe, S. A., Spence, J. C., Friedman, A., et al. (2015). Systematic review of sedentary behavior and cognitive development in early childhood. *Prev. Med.* 78, 115–122. doi: 10.1016/j.yjmed.2015.07.016
- Castebon, K., and Andreyeva, T. (2012). Obesity and motor skills among 4 to 6-year-old children in the United States: nationally-representative surveys. *BMC Pediatr.* 12:28. doi: 10.1186/1471-2431-12-28
- Cole, T. J., Bellizzi, M. C., Flegal, K. M., and Dietz, W. H. (2000). Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 320, 1240–1243. doi: 10.1136/bmj.320.7244.1240
- Cornoldi, C., Miato, L., Molin, A., and Poli, S. (2009). *PRCR-2/2009. Prove di prerequisito per la diagnosi delle difficoltà di lettura e scrittura*. Firenze: Giunti Organizzazioni Speciali.
- Czenczek-Lewandowska, E., Grzegorzczak, J., and Mazur, A. (2018). Physical activity in children and adolescents with type 1 diabetes and contemporary methods of its assessment. *Pediatr. Endocrinol. Diabetes Metab.* 24, 179–184. doi: 10.5114/pedm.2018.83364
- Deli, E., Bakle, I., and Zachopoulou, E. (2006). Implementing intervention movement programs for kindergarten children. *J. Early Childhood Res.* 4, 5–18. doi: 10.1177/1476718X06059785
- D'Hondt, E., Deforche, B., De Bourdeaudhuij, I., and Lenoir, M. (2009). Relationship between motor skill and body mass index in 5- to 10-year-old children. *Adapt. Phys. Activ. Q* 26, 21–37. doi: 10.1123/apaq.26.1.21
- Donnelly, J. E., Hillman, C. H., Castelli, D., Etnier, J. L., Lee, S., Tomporowski, P., et al. (2016). Physical activity, fitness, cognitive function, and academic achievement in children: a systematic review. *Med. Sci. Sports Exerc.* 48, 1197–1222. doi: 10.1249/MSS.0000000000000901
- DuBose, K. D., Gross McMillan, A., Wood, A. P., and Sisson, S. B. (2018). Joint relationship between physical activity, weight status, and motor skills in children aged 3 to 10 years. *Percept. Mot. Skills* 125, 478–492. doi: 10.1177/0031512518767008
- Eichinger, M., Schneider, S., and De Bock, F. (2017). Subjectively and objectively assessed social and physical environmental correlates of preschoolers' accelerometer-based physical activity. *Int. J. Behav. Nutr. Phys. Act.* 14:153. doi: 10.1186/s12966-017-0577-9
- Engel, A. C., Broderick, C. R., van Doorn, N., Hardy, L. L., and Parmenter, B. J. (2018). Exploring the relationship between fundamental motor skill interventions and physical activity levels in children: a systematic review and meta-analysis. *Sports Med.* 48, 1845–1857. doi: 10.1007/s40279-018-0923-3
- Figueroa, R., and An, R. (2017). Motor skill competence and physical activity in preschoolers: a review. *Matern. Child Health J.* 21, 136–146. doi: 10.1007/s10995-016-2102-1
- Fisher, A., Reilly, J. J., Kelly, L. A., Montgomery, C., Williamson, A., Paton, J. Y., et al. (2005). Fundamental movement skills and habitual physical activity in young children. *Med. Sci. Sports Exerc.* 37, 684–688. doi: 10.1249/01.MSS.0000159138.48107.7D
- Giustino, V., Parroco, A. M., Gennaro, A., Musumeci, G., Palma, A., and Battaglia, G. (2020). Physical activity levels and related energy expenditure during COVID-19 quarantine among the sicilian active population: a cross-sectional online survey study. *Sustainability* 12:4356. doi: 10.3390/su12114356
- Goodway, J. D., Crowe, H., and Ward, P. (2003). Effects of motor skill instruction on fundamental motor skill development. *Adapt. Phys. Act. Q* 20, 298–314. doi: 10.1123/apaq.20.3.298
- Gu, X. (2016). Fundamental motor skill, physical activity, and sedentary behavior in socioeconomically disadvantaged kindergarteners. *Psychol. Health Med.* 21, 871–881. doi: 10.1080/13548506.2015.1125007
- Han, A., Fu, A., Cobley, S., and Sanders, R. H. (2018). Effectiveness of exercise intervention on improving fundamental movement skills and motor coordination in overweight/obese children and adolescents: a systematic review. *J. Sci. Med. Sport* 21, 89–102. doi: 10.1016/j.jsams.2017.07.001
- Han, J. C., Lawlor, D. A., and Kimm, S. Y. (2010). Childhood obesity. *Lancet* 375, 1737–1748. doi: 10.1016/S0140-6736(10)60171-7
- Hardy, L. L., King, L., Kelly, B., Farrell, L., and Howlett, S. (2010). Munch and move: evaluation of a preschool healthy eating and movement skill program. *Int. J. Behav. Nutr. Phys. Act.* 7:80. doi: 10.1186/1479-5868-7-80
- Hills, A. P., Andersson, L. B., and Byrne, N. M. (2011). Physical activity and obesity in children. *Br. J. Sports Med.* 45, 866–870. doi: 10.1136/bjsports-2011-090199
- Hoza, B., Shoulberg, E. K., Tompkins, C. L., Meyer, L. E., Martin, C. P., Krasner, A., et al. (2020). Meeting a physical activity guideline in preschool and school readiness: a program evaluation. *Child Psychiatry Hum. Dev.* doi: 10.1007/s10578-020-01055-9. [Epub ahead of print].
- Jarraya, S., Wagner, M., Jarraya, M., and Engel, F. A. (2019). 12 Weeks of Kindergarten-based yoga practice increases visual attention, visual-motor precision and decreases behavior of inattention and hyperactivity in 5-year-old children. *Front. Psychol.* 10:796. doi: 10.3389/fpsyg.2019.00796
- Kim, C., and Lee, K. (2016). The relationship between fundamental movement skills and body mass index in Korean preschool children. *Eur. Early Child. Educ. Res. J.* 24, 928–935. doi: 10.1080/1350293X.2016.1239326
- Kimbrow, R. T., Brooks-Gunn, J., and McLanahan, S. (2011). Young children in urban areas: links among neighborhood characteristics, weight status, outdoor play, and television watching. *Soc. Sci. Med.* 72, 668–676. doi: 10.1016/j.socscimed.2010.12.015
- Kippe, K. O., and Lagestad, P. A. (2018). Kindergarten: producer or reducer of inequality regarding physical activity levels of preschool children. *Front. Public Health* 6:361. doi: 10.3389/fpubh.2018.00361
- Kirk, S. M., and Kirk, E. P. (2016). Sixty minutes of physical activity per day included within preschool academic lessons improves early literacy. *J. Sch. Health* 86, 155–163. doi: 10.1111/josh.12363
- Kirk, S. M., Vizcarra, C. R., Looney, E. C., and Kirk, E. P. (2014). Using physical activity to teach academic content: a study of the effects on literacy in head start preschoolers. *Early Childhood Educ. J.* 42, 181–189. doi: 10.1007/s10643-013-0596-3
- Kobel, S., Wartha, O., Lammle, C., Dreyhaupt, J., and Steinacker, J. M. (2019). Intervention effects of a kindergarten-based health promotion programme on obesity related behavioural outcomes and BMI percentiles. *Prev. Med. Rep.* 15:100931. doi: 10.1016/j.pmedr.2019.100931
- Konstabel, K., Veidebaum, T., Verbestel, V., Moreno, L. A., Bammann, K., Tornaritis, M., et al. (2014). Objectively measured physical activity in European children: the IDEFICS study. *Int. J. Obes.* 38, S135–S143. doi: 10.1038/ijo.2014.144
- Lai, S. K., Costigan, S. A., Morgan, P. J., Lubans, D. R., Stodden, D. F., Salmon, J., et al. (2014). Do school-based interventions focusing on physical activity, fitness, or fundamental movement skill competency produce a sustained impact in these outcomes in children and adolescents? A systematic review of follow-up studies. *Sports Med.* 44, 67–79. doi: 10.1007/s40279-013-0099-9
- Liang, J., Matheson, B. E., Kaye, W. H., and Boutelle, K. N. (2014). Neurocognitive correlates of obesity and obesity-related behaviors in children and adolescents. *Int. J. Obes.* 38, 494–506. doi: 10.1038/ijo.2013.142
- Logan, S. W., Robinson, L. E., Wilson, A. E., and Lucas, W. A. (2012). Getting the fundamentals of movement: a meta-analysis of the effectiveness of motor skill interventions in children. *Child Care Health Dev.* 38, 305–315. doi: 10.1111/j.1365-2214.2011.01307.x

- Logan, S. W., Scrabis-Fletcher, K., Modlesky, C., and Getchell, N. (2011). The relationship between motor skill proficiency and body mass index in preschool children. *Res. Q. Exerc. Sport* 82, 442–448. doi: 10.1080/02701367.2011.10599776
- Lohman, T. G., Roche, A. F., and Martorell, R. (1988). *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics Books.
- Lonigan, C. J., and Shanahan, T. (2010). Developing early literacy skills: things we know we know and things we know we don't know. *Educ. Res.* 39, 340–346. doi: 10.3102/0013189X10369832
- Lopes, V. P., Stodden, D. F., Bianchi, M. M., Maia, J. A., and Rodrigues, L. P. (2012). Correlation between BMI and motor coordination in children. *J. Sci. Med. Sport* 15, 38–43. doi: 10.1016/j.jsams.2011.07.005
- Loras, H. (2020). The effects of physical education on motor competence in children and adolescents: a systematic review and meta-analysis. *Sports* 8:88. doi: 10.3390/sports8060088
- Maeng, H., Webster, E. K., Pitchford, E. A., and Ulrich, D. A. (2017). Inter- and intrarater reliabilities of the test of gross motor development-third edition among experienced TGMD-2 raters. *Adapt. Phys. Activ. Q* 34, 442–455. doi: 10.1123/apaq.2016-0026
- Mavilidi, M., Okely, A. D., Chandler, P., Cliff, D. P., and Paas, F. (2015). Effects of integrated physical exercises and gestures on preschool children's foreign language vocabulary learning. *Educ. Psychol. Rev.* 27, 413–426. doi: 10.1007/s10648-015-9337-z
- Mavilidi, M., Okely, A. D., Chandler, P., and Paas, F. (2017). Effects of integrating physical activities into a science lesson on preschool children's learning and enjoyment. *Appl. Cogn. Psychol.* 31, 281–290. doi: 10.1002/acp.3325
- Morano, M., Colella, D., and Caroli, M. (2011). Gross motor skill performance in a sample of overweight and non-overweight preschool children. *Int. J. Pediatr. Obes.* 6, 42–46. doi: 10.3109/17477166.2011.613665
- Mülazimoglu Ballı, O. (2016). Motor proficiency and body mass index of preschool children: in relation to socioeconomic status. *J. Educ. Train Stud.* 4, 237–243. doi: 10.11114/jets.v4i4.1302
- Nielsen, G., Mygind, E., Bolling, M., Otte, C. R., Schneller, M. B., Schipperijn, J., et al. (2016). A quasi-experimental cross-disciplinary evaluation of the impacts of education outside the classroom on pupils' physical activity, well-being and learning: the TEACHOUT study protocol. *BMC Public Health* 16:1117. doi: 10.1186/s12889-016-3780-8
- Nilsen, A. K. O., Anderssen, S. A., Johannessen, K., Aadland, K. N., Ylvisaker, E., Loftesnes, J. M., et al. (2020a). Bi-directional prospective associations between objectively measured physical activity and fundamental motor skills in children: a two-year follow-up. *Int. J. Behav. Nutr. Phys. Act.* 17:1. doi: 10.1186/s12966-019-0902-6
- Nilsen, A. K. O., Anderssen, S. A., Loftesnes, J. M., Johannessen, K., Ylvisaker, E., and Aadland, E. (2020b). The multivariate physical activity signature associated with fundamental motor skills in preschoolers. *J. Sports Sci.* 38, 264–272. doi: 10.1080/02640414.2019.1694128
- Oberer, N., Gashaj, V., and Roebers, C. M. (2017). Motor skills in kindergarten: internal structure, cognitive correlates and relationships to background variables. *Hum. Mov. Sci.* 52, 170–180. doi: 10.1016/j.humov.2017.02.002
- Obradovic Salcin, L., Karin, Z., Miljanovic Damjanovic, V., Ostojic, M., Vrdoljak, A., Gilic, B., et al. (2019). Physical activity, body mass, and adherence to the mediterranean diet in preschool children: a cross-sectional analysis in the split-dalmatia county (Croatia). *Int. J. Environ. Res. Public Health* 16:3237. doi: 10.3390/ijerph16183237
- Orton, J., Spittle, A., Doyle, L., Anderson, P., and Boyd, R. (2009). Do early intervention programmes improve cognitive and motor outcomes for preterm infants after discharge? A systematic review. *Dev. Med. Child Neurol.* 51, 851–859. doi: 10.1111/j.1469-8749.2009.03414.x
- Osinski, W., and Kantanista, A. (2017). Physical activity in the therapy of overweight and obesity in children and adolescents. Needs and recommendations for intervention programs. *Dev. Period Med.* 21, 224–234.
- Pagani, L. S., Fitzpatrick, C., Barnett, T. A., and Dubow, E. (2010). Prospective associations between early childhood television exposure and academic, psychosocial, and physical well-being by middle childhood. *Arch. Pediatr. Adolesc. Med.* 164, 425–431. doi: 10.1001/archpediatrics.2010.50
- Palmer, K. K., Chinn, K. M., and Robinson, L. E. (2019). The effect of the CHAMP intervention on fundamental motor skills and outdoor physical activity in preschoolers. *J. Sport Health Sci.* 8, 98–105. doi: 10.1016/j.jsbs.2018.12.003
- Pinto, G., Bigozzi, L., Tarchi, C., Vezzani, C., and Accorti Gamannossi, B. (2016). Predicting reading, spelling, and mathematical skills: a longitudinal study from kindergarten through first grade. *Psychol. Rep.* 118, 413–440. doi: 10.1177/0033294116633357
- Popovic, B., Cvetkovic, M., Macak, D., Scepanovic, T., Cokorilo, N., Belic, A., et al. (2020). Nine months of a structured multisport program improve physical fitness in preschool children: a quasi-experimental study. *Int. J. Environ. Res. Public Health* 17:4935. doi: 10.3390/ijerph17144935
- Puranik, C. S., and Lonigan, C. J. (2011). From scribbles to scrabble: preschool children's developing knowledge of written language. *Read. Writ.* 24, 567–589. doi: 10.1007/s11145-009-9220-8
- Reilly, J. J., Hughes, A. R., Gillespie, J., Malden, S., and Martin, A. (2019). Physical activity interventions in early life aimed at reducing later risk of obesity and related non-communicable diseases: a rapid review of systematic reviews. *Obes. Rev.* 20, 61–73. doi: 10.1111/obr.12773
- Rhemtulla, M., and Tucker-Drob, E. M. (2011). Correlated longitudinal changes across linguistic, achievement, and psychomotor domains in early childhood: evidence for a global dimension of development. *Dev. Sci.* 14, 1245–1254. doi: 10.1111/j.1467-7687.2011.01071.x
- Rodriguez-Ayllon, M., Cadenas-Sanchez, C., Estevez-Lopez, F., Munoz, N. E., Mora-Gonzalez, J., Migueles, J. H., et al. (2019). Role of physical activity and sedentary behavior in the mental health of preschoolers, children and adolescents: a systematic review and meta-analysis. *Sports Med.* 49, 1383–1410. doi: 10.1007/s40279-019-01099-5
- Roscoe, C. M. P., James, R. S., and Duncan, M. J. (2019). Accelerometer-based physical activity levels, fundamental movement skills and weight status in British preschool children from a deprived area. *Eur. J. Pediatr.* 178, 1043–1052. doi: 10.1007/s00431-019-03390-z
- Saraiva, L., Rodrigues, L. P., Cordovil, R., and Barreiros, J. (2013). Influence of age, sex and somatic variables on the motor performance of preschool children. *Ann. Hum. Biol.* 40, 444–450. doi: 10.3109/03014460.2013.802012
- Siakhkouhian, M., Mahmoodi, H., and Salehi, M. (2011). Relationship between fundamental movement skills and body mass index in 7-to-8 year-old children. *World Appl. Sci. J.* 15, 1354–1360.
- Stodden, D. F., Goodway, J. D., Langendorfer, S. J., Roberton, M. A., Rudisill, M. E., Garcia, C., et al. (2008). A developmental perspective on the role of motor skill competence in physical activity: an emergent relationship. *Quest* 60, 290–306. doi: 10.1080/00336297.2008.10483582
- Tabacchi, G., Battaglia, G., Alesi, M., Paoli, A., Palma, A., and Bellafiore, M. (2020a). Food literacy predictors and associations with physical and emergent literacy in pre-schoolers: results from the training-to-health project. *Public Health Nutr.* 23, 356–365. doi: 10.1017/S1368980019002404
- Tabacchi, G., Battaglia, G., Messina, G., Paoli, A., Palma, A., and Bellafiore, M. (2020b). Validity and internal consistency of the preschool-FLAT, a new tool for the assessment of food literacy in young children from the training-to-health project. *Int. J. Environ. Res. Public Health* 17:2759. doi: 10.3390/ijerph17082759
- Toussaint, N., Streppel, M. T., Mul, S., Fukkink, R. G., Weijs, P. J. M., and Janssen, M. (2020). The effects of the PLAYTOD program on children's physical activity at preschool playgrounds in a deprived Urban area: a randomized controlled trial. *Int. J. Environ. Res. Public Health* 17:329. doi: 10.3390/ijerph17010329
- Tsuda, E., Goodway, J. D., Famelia, R., and Brian, A. (2020). Relationship between fundamental motor skill competence, perceived physical competence and free-play physical activity in children. *Res. Q. Exerc. Sport* 91, 55–63. doi: 10.1080/02701367.2019.1646851
- Tucker, P. (2008). The physical activity levels of preschool-aged children: a systematic review. *Early Child. Res. Q* 23, 547–558. doi: 10.1016/j.ecresq.2008.08.005
- Ulrich, D. A. (2003). *Test TGM-Valutazione Delle Abilita' Grosse-Motorie*. Trento: Erickson.
- Van Capelle, A., Broderick, C. R., van Doorn, N., R, E.W., and Parmenter, B. J. (2017). Interventions to improve fundamental motor skills in pre-school aged children: a systematic review and meta-analysis. *J. Sci. Med. Sport* 20, 658–666. doi: 10.1016/j.jsams.2016.11.008

- Voss, C., and Harris, K. C. (2017). Physical activity evaluation in children with congenital heart disease. *Heart* 103, 1408–1412. doi: 10.1136/heartjnl-2017-311340
- Xin, F., Chen, S. T., Clark, C., Hong, J. T., Liu, Y., and Cai, Y. J. (2020). Relationship between fundamental movement skills and physical activity in preschool-aged children: a systematic review. *Int. J. Environ. Res. Public Health* 17:3566. doi: 10.3390/ijerph17103566
- Zeng, N., Ayyub, M., Sun, H., Wen, X., Xiang, P., and Gao, Z. (2017). Effects of physical activity on motor skills and cognitive development in early childhood: a systematic review. *Biomed Res. Int.* 2017:2760716. doi: 10.1155/2017/2760716

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Battaglia, Giustino, Tabacchi, Alesi, Galassi, Modica, Palma and Bellafore. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.